

Opinion

Apoptosis induced by *Pseudomonas aeruginosa*: a lonely killer?

Alexis Broquet^{1*} and Karim Asehnoune^{1,2}

¹Faculté de Médecine, Laboratoire UPRES EA 3826, Université de Nantes, Nantes, France.

²Pôle Anesthésie Réanimations, Service d'Anesthésie Réanimation Chirurgicale, Hôtel Dieu, CHU Nantes, Nantes, France.

Apoptosis is a fundamental biological process allowing tissue homeostasis through the regulation of cell populations by eliminating unnecessary elements. During infection, pathogens have evolved to take advantage of this process for their own and are able to induce the apoptosis of cells, i.e. immune cells by the host itself.

Pseudomonas aeruginosa is one of the most studied opportunistic bacteria due to its significant involvement worldwide in pneumonia, corneal infections and wound burns. Several research groups have pointed out the ability of these bacteria to interfere and/or evade host immune system by inducing apoptosis of the targeted cells. In May 2014, looking up '*Pseudomonas aeruginosa*' and 'apoptosis' keywords in PubMed search engine retrieve more than 300 hits. *Pseudomonas aeruginosa* seems to induce apoptosis through direct interaction with the host cells (the most studied system being the type-III secretion system: T3SS) or through secreting factors such as pyocyanin.

T3SS, the most well-studied virulence apparatus of *P. aeruginosa* is composed of a needle complex through which exoenzymes are injected into the host cells (Galle *et al.*, 2012). Recently, Beyaert's laboratory described an exotoxin-independent function of the T3SS in the killing of macrophages in an acute lung infection model (Galle *et al.*, 2012). Although, T3SS is a major virulence system, it is not fully required for the bacteria to display virulence as T3SS negative strains are shown to exhibit significant virulence (example of Elsen's paper). In short, a wide variety of *P. aeruginosa* virulence factors are involved in inducing apoptosis by several distinct mechanisms, from the activation of the mitochondrial pathway,

the generation of reactive oxygen species to the activation of the caspase pathways (Table 1).

The ability of *P. aeruginosa* to induce apoptosis in various *in vitro* model of infection (macrophages, neutrophils, epithelial cells . . .) or *in vivo* models such as lungs, cornea and burn wounds infections is not mediated by a single bacterial cell but rather by a multicellular population of *P. aeruginosa*. Members of such population interact with each other through a number of chemical signals known globally on for quorum sensing (QS). Quoting Rutherford and Bassler (2012), 'Quorum sensing is a bacterial cell-cell communication process that involves the production, detection, and response to extracellular signaling molecules called autoinducers'. QS molecules were shown to regulate virulence factors such as toxins, exotoxin A, pyocyanin, . . . and so *in fine* apoptosis (Rutherford and Bassler, 2012). The best described QS signalling systems in *P. aeruginosa* are the N-acyl homoserine lactones systems Las and Rhl. The Las system produces and responds to N-oxododecanoyl homoserine lactone and the Rhl system to N-butanoyl homoserine lactone respectively. Las system is known to control the production of various virulence factors involved in host cell damages such as exotoxin A (Jones *et al.*, 1993). On the other hand, Rhl system was described to repress the expression of genes responsible for the assembly and function of the T3SS (Bleves *et al.*, 2005).

Last but not least, QS molecule such as 3-oxododecanoyl-L-homoserine lactone (3-oxo-C12-HSL) itself has been shown to induce apoptosis. Several studies have demonstrated that incubation of different cell lines with 3-oxo-C12-HSL molecule resulted in the induction of apoptosis involving calcium signalling, the mitochondrial pathway and caspase activations (Table 2). Interestingly, N-butanoyl-L-homoserine lactone (known as C4-HSL, the second major QS molecule in *P. aeruginosa*) harbouring a shorter fatty acid chain has not been shown to induce apoptosis compared with 3-oxo-C12-HSL (Tateda *et al.*, 2003; Holban *et al.*, 2014).

Knowing that *P. aeruginosa* is able to induce apoptosis through its QS systems molecule, studies focusing on apoptosis induction should be considered with the context of QS signalling. Particularly, QS considerations should be taken into account when comparing studies using

Received 11 June, 2014; accepted 13 June, 2014. *For correspondence. E-mail alexis.broquet@univ-nantes.fr; Tel. +33 (0)2 40 41 28 38.

doi:10.1111/1751-7915.12144

Funding Information No funding information provided.

Table 1. Virulence factors inducing apoptosis in targeted host cells (non-exhaustive list).

Virulence factors	Model used	Apoptosis pathway	Reference
Type-III secretion system (T3SS)			
Injector/needle complex	Macrophages/neutrophils	Caspase 3	Galle <i>et al.</i> , 2012
Exoenzymes (ExoS, T, U, Y)	Macrophages	Caspase 3	Galle <i>et al.</i> , 2008
	Epithelial cells/fibroblasts	Mitochondrial pathway, caspase 3	Shafikhani <i>et al.</i> , 2008
Secreted virulence factors			
Pyocyanin	Neutrophils	Reactive oxygen intermediate, cAMP	Usher <i>et al.</i> , 2002
Exotoxin A	MEFs	Bak pathway	Du <i>et al.</i> , 2010
Protease	Macrophages	Caspase 3	Zhang <i>et al.</i> , 2003
ExlA	HUVECs endothelial cells	Unknown	Elsen <i>et al.</i> , 2014

cAMP, cyclic adenosine monophosphate; HUVECs, human umbilical veins; MEFs, mouse embryonic fibroblasts.

Table 2. Apoptosis pathways activated by 3-oxo-C12-HSL (non-exhaustive list).

Quorum-sensing molecule	Abbreviation used in the study	Model used	Apoptosis pathway involved	Reference
N-3-(oxododecanoyl)-l-homoserine lactone	3-oxo-C12-HSL	Macrophages/neutrophils	Caspase 3/8	Tateda <i>et al.</i> , 2003
	OdDHL	Breast cancer cell lines	JAK/STAT pathway	Li <i>et al.</i> , 2004
	OdDHL	Jurkat cell line	Mitochondrial pathway	Jacobi <i>et al.</i> , 2009
	C12	Airway epithelial cells	Cytochrome <i>c</i> , caspases 3/7, 8 and 9	Schwarzer <i>et al.</i> , 2012

different multiplicity of infection or bacteria preparation protocols, processes that influence QS molecules concentration and/or bacteria population numbers.

Conflict of interest

None declared.

References

- Bleves, S., Soscia, C., Nogueira-Orlandi, P., Lazdunski, A., and Filloux, A. (2005) Quorum sensing negatively controls type III secretion regulon expression in *Pseudomonas aeruginosa* PAO1. *J Bacteriol* **187**: 3898–3902.
- Du, X., Youle, R.J., FitzGerald, D.J., and Pastan, I. (2010) *Pseudomonas* exotoxin A-mediated apoptosis is Bak dependent and preceded by the degradation of Mcl-1. *Mol Cell Biol* **30**: 3444–3452.
- Elsen, S., Huber, P., Bouillot, S., Couté, Y., Fournier, P., Dubois, Y., *et al.* (2014) A type III secretion negative clinical strain of *Pseudomonas aeruginosa* employs a two-partner secreted exolysin to induce hemorrhagic pneumonia. *Cell Host Microbe* **15**: 164–176.
- Galle, M., Schotte, P., Haegman, M., Wullaert, A., Yang, H.J., Jin, S., and Beyaert, R. (2008) The *Pseudomonas aeruginosa* type III secretion system plays a dual role in the regulation of caspase-1 mediated IL-1 β maturation. *J Cell Mol Med* **12** (5A): 1767–1776.
- Galle, M., Carpentier, I., and Beyaert, R. (2012) Structure and function of the Type III secretion system of *Pseudomonas aeruginosa*. *Curr Protein Pept Sci* **13**: 831–842.
- Galle, M., Jin, S., Bogaert, P., Haegman, M., Vandenabeele, P., and Beyaert, R. (2012) The *Pseudomonas aeruginosa* type III secretion system has an exotoxin S/T/Y independent pathogenic role during acute lung infection. *PLoS ONE* **7**: e41547.
- Holban, A.M., Bleotu, C., Chifiriuc, M.C., Bezirtzoglou, E., and Lazar, V. (2014) Role of *Pseudomonas aeruginosa* quorum sensing (QS) molecules on the viability and cytokine profile of human mesenchymal stem cells. *Virulence* **5**: 303–310.
- Jacobi, C.A., Schiffner, F., Henkel, M., Waibel, M., Stork, B., Daubrawa, M., *et al.* (2009) Effects of bacterial N-acyl homoserine lactones on human Jurkat T lymphocytes—OdDHL induces apoptosis via the mitochondrial pathway. *Int J Med Microbiol* **299**: 509–519.
- Jones, S., Yu, B., Bainton, N.J., Birdsall, M., Bycroft, B.W., Chhabra, S.R., *et al.* (1993) The lux autoinducer regulates the production of exoenzyme virulence determinants in *Erwinia carotovora* and *Pseudomonas aeruginosa*. *EMBO J* **12**: 2477–2482.
- Li, L., Hooi, D., Chhabra, S.R., Pritchard, D., and Shaw, P.E. (2004) Bacterial N-acylhomoserine lactone-induced apoptosis in breast carcinoma cells correlated with down-modulation of STAT3. *Oncogene* **23**: 4894–4902.
- Rutherford, S.T., and Bassler, B.L. (2012) Bacterial quorum sensing: its role in virulence and possibilities for its control. *Cold Spring Harb Perspect Med* **2**: a012427.
- Schwarzer, C., Fu, Z., Patanwala, M., Hum, L., Lopez-Guzman, M., Illek, B., *et al.* (2012) *Pseudomonas aeruginosa* biofilm-associated homoserine lactone C12 rapidly activates apoptosis in airway epithelia. *Cell Microbiol* **14**: 698–709.
- Shafikhani, S.H., Morales, C., and Engel, J. (2008) The *Pseudomonas aeruginosa* type III secreted toxin ExoT is necessary and sufficient to induce apoptosis in epithelial cells. *Cell Microbiol* **10**: 994–1007.
- Tateda, K., Ishii, Y., Horikawa, M., Matsumoto, T., Miyairi, S., Pechere, J.C., *et al.* (2003) The *Pseudomonas aeruginosa*

autoinducer N-3-oxododecanoyl homoserine lactone accelerates apoptosis in macrophages and neutrophils. *Infect Immun* **71**: 5785–5793.

Usher, L.R., Lawson, R.A., Geary, I., Taylor, C.J., Bingle, C.D., Taylor, G.W., and Whyte, M.K. (2002) Induction of neutrophil apoptosis by the *Pseudomonas aeruginosa*

exotoxin pyocyanin: a potential mechanism of persistent infection. *J Immunol* **168**: 1861–1868.

Zhang, J., Takayama, H., Matsuba, T., Jiang, R., and Tanaka, Y. (2003) Induction of apoptosis in macrophage cell line, J774, by the cell-free supernatant from *Pseudomonas aeruginosa*. *Microbiol Immunol* **47**: 199–206.